



Eigenvalues of graphs and a simple proof of a theorem of Greenberg

Sebastian M. Cioabă^{*,1}

Department of Mathematics, Queen's University at Kingston, Ont., Canada K7L 3N6

Received 16 March 2005; accepted 21 December 2005

Available online 9 February 2006

Submitted by R.A. Brualdi

Abstract

In his Ph.D. thesis, Greenberg proved that if $\rho(\tilde{X})$ is the spectral radius of the universal cover \tilde{X} of a finite graph X , then for each $\epsilon > 0$, a positive proportion (depending only on \tilde{X} and ϵ) of the eigenvalues of X have *absolute value* at least $\rho(\tilde{X}) - \epsilon$. In this paper, we show that the same result holds true if we remove *absolute* from the previous result. We also prove an analogue result for the smallest eigenvalues of X .
© 2006 Elsevier Inc. All rights reserved.

AMS classification: 05C50; 15A18

Keywords: Spectral radius; Universal cover; Eigenvalues

1. Preliminaries

Our graph theoretic notation is standard, see [18]. The graphs discussed in this paper are simple and connected unless stated otherwise. For a graph X , we denote by $\lambda_1(X) \geq \lambda_2(X) \geq \dots \geq \lambda_n(X)$ the eigenvalues of the adjacency matrix of X .

Serre has proved the following theorem (see [4,5,11,17]) using Chebyshev polynomials. The simplest self-contained proof of this theorem is given in [4] and it is fairly involved. See [2,3] for a simple proof of Serre's theorem and other related results.

* Present address: Department of Mathematics, Dept 0112, University of California at San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0112, USA.

E-mail address: sebi.cioaba@gmail.com

¹ Research partially supported by an Ontario Graduate Scholarship.

Theorem 1 (Serre). *For each $\epsilon > 0$, there exists a positive constant $c = c(\epsilon, k)$ such that for any k -regular graph X , the number of eigenvalues λ of X with $\lambda \geq (2 - \epsilon)\sqrt{k - 1}$ is at least $c|X|$.*

Serre’s theorem is a generalization of the asymptotic Alon and Boppana theorem, see [1,6,16] for more details.

Theorem 2. *If $(X_n)_n$ is an infinite sequence of k -regular graphs, then $\liminf \lambda_2(X_n) \geq 2\sqrt{k - 1}$.*

Theorem 1 is also related to a result obtained by Greenberg in [9] whose proof has not appeared to our knowledge in any journal as of yet. Greenberg’s result is cited in many places, [14,15] for example. For a graph X , we denote by $\rho(X)$ its spectral radius and by $\mathcal{C}(X)$ the family of all finite graphs that are covered by X . In Section 2, we describe these notions in more detail.

Theorem 3 (Greenberg). *Let X be a connected, infinite graph with finite maximum degree. Given $\epsilon > 0$, there exists $c = c(X, \epsilon) > 0$, such that for every $Y \in \mathcal{C}(X)$,*

$$|\{\lambda \in \text{spectrum of } Y : |\lambda| \geq \rho(X) - \epsilon\}| \geq c|V(Y)|. \tag{1}$$

In Section 3, we present a proof of Theorem 3. Note that Theorem 3 implies a weaker form of Serre’s theorem. This is because if X is the infinite k -regular tree, then $\rho(X) = 2\sqrt{k - 1}$. Thus, if Y is a finite k -regular graph we obtain that for each $\epsilon > 0$, a positive proportion (that depends only on ϵ and k) of the eigenvalues of Y have *absolute value* at least $(2 - \epsilon)\sqrt{k - 1}$. This is slightly weaker than Theorem 1.

In Section 4, we present a slight improvement of Greenberg’s theorem. We prove that given X and $\epsilon > 0$, there is a positive $c = c(X, \epsilon)$ such that $|\{\lambda \in \text{spectrum of } Y : \lambda \geq \rho(X) - \epsilon\}| \geq c|V(Y)|$ for each finite graph Y covered by X . We also prove a similar result regarding the smallest eigenvalues of general (not necessarily regular) graphs.

2. Graphs and coverings

If X is a connected graph (not necessarily finite) such that the maximum degree of X is finite, let $l^2(X)$ denote the space of functions $f : V(X) \rightarrow \mathbb{R}$ with $\sum_{x \in V(X)} |f(x)|^2 < +\infty$. Let $\delta : l^2(X) \rightarrow l^2(X)$ be the adjacency operator of X , i.e., $(\delta f)(x) = \sum_{y: x \in E(X)} f(y)$. If $x \in V(X)$, let $t_s(x)$ denote the number of closed walks of length s that start at x . Denote by $\rho(X)$ the *spectral radius* of X :

$$\rho(X) = \sup\{|\lambda| : \lambda \in \text{spectrum of } \delta\}.$$

Lemma 4. *Let X be a connected graph. Then $\limsup_{s \rightarrow +\infty} \sqrt[s]{t_s(x)}$ is independent on the vertex $x \in V(X)$.*

Proof. Since X is connected, it is enough to prove that

$$\limsup_{s \rightarrow +\infty} \sqrt[s]{t_s(x)} = \limsup_{s \rightarrow +\infty} \sqrt[s]{t_s(y)}$$

for any adjacent vertices x and y . The previous assertion will follow easily from the fact that

$$t_{s+2}(y) \geq t_s(x) \geq t_{s-2}(y)$$

for each $s \geq 2$. \square

It is well known (cf. [13, Chapter 4]) that

$$\rho(X) = \limsup_{s \rightarrow +\infty} \sqrt[s]{t_s(x)}. \tag{2}$$

Given two graphs X_1 and X_2 , a *homomorphism* from X_1 to X_2 is a function $f : V(X_1) \rightarrow V(X_2)$ such that $xy \in E(X_1)$ implies $f(x)f(y) \in E(X_2)$ for each $x, y \in V(X_1)$. If f is bijective and f and f^{-1} are both homomorphisms, then f is called an *isomorphism* from X_1 to X_2 . An isomorphism from a graph X to itself is called an *automorphism* of X . The automorphisms of X form a group, called the *automorphism group* of X that we denote by $\text{Aut}(X)$. If x is a vertex of X , then the *automorphism orbit* of x is $\text{Orb}(x) = \{y \in V(X) : \exists f \in \text{Aut}(X) \text{ such that } f(x) = y\}$. If x is a vertex in the graph X , then we denote by $N_X(x)$ the set of neighbours of x in X .

If X_1 and X_2 are two graphs, a homomorphism $\pi : V(X_1) \rightarrow V(X_2)$ is called a *cover map* if it is surjective and for each $x \in V(X_1)$, π induces an isomorphism from $N_{X_1}(x)$ to $N_{X_2}(\pi(x))$. It follows from (2) that if $\pi : V(X_1) \rightarrow V(X_2)$ is a cover map, then $\rho(X_1) \leq \rho(X_2)$. If π is a finite cover, then $\rho(X_1) = \rho(X_2)$. Denote by $\mathcal{C}(X)$ the family of finite graphs covered by X .

Using a result of Leighton [12], the next theorem is also proved by Greenberg [9] (see also [14]).

Theorem 5. *Let X be a connected graph with finite maximum degree. Then for each X_1 and X_2 in $\mathcal{C}(X)$, $\rho(X_1) = \rho(X_2)$. This common value is denoted by $\chi(X)$.*

For a finite graph Z , its *universal cover* \tilde{Z} is the graph with the property that for any graph Y with a cover map $\pi : V(Y) \rightarrow V(Z)$, there exists a cover map $\pi' : V(\tilde{Z}) \rightarrow V(Y)$. The universal cover of any finite graph is an infinite tree. For example, the universal cover of a k -regular graph is the infinite k -regular tree. However, not every infinite tree X covers a finite graph. It is easy to see that a necessary condition for covering a finite graph is that $\text{Aut}(X)$ has finitely many orbits. See [14] for more details on the universal covers of finite graphs.

3. A proof of Greenberg’s theorem

In his Ph.D. thesis [9], Greenberg proved Theorem 3. This result is also cited in [14,15, Theorem 2.3], but it seems that no proof of it exists in the literature other than in Greenberg’s thesis. The proof given below is a simplified version of the original proof.

Proof of Theorem 3. Let $\epsilon > 0$ and $Y \in \mathcal{C}(X)$. Because X has finitely many automorphism orbits, it follows that there exists $r_0 = r(X, \epsilon)$ such that

$$t_{2r}(y) \geq \left(\rho(X) - \frac{\epsilon}{2}\right)^{2r}$$

for each vertex $y \in V(Y)$ and $r \geq r_0$. Let c be the proportion of eigenvalues of Y that have absolute value $\geq \rho(X) - \epsilon$. Using the previous inequality, we obtain

$$\left(\rho(X) - \frac{\epsilon}{2}\right)^{2r} \leq \min_{y \in V(Y)} t_{2r}(y) \leq \frac{\text{tr}(A^{2r}(Y))}{|V(Y)|} \leq c\chi^{2r}(X) + (1 - c)(\rho(X) - \epsilon)^{2r}$$

for each $r \geq r_0$. This implies

$$c \geq \frac{(\rho(X) - \frac{\epsilon}{2})^{2r} - (\rho(X) - \epsilon)^{2r}}{\chi^{2r}(X) - (\rho(X) - \frac{\epsilon}{2})^{2r}}$$

for each $r \geq r_0$. Letting $r = r_0$, this proves the theorem. \square

4. An slight improvement of Greenberg’s theorem

In this section, we present a slight improvement of Theorem 3. Our proof is similar to the previous one and we obtain the required estimate for the largest eigenvalues by shifting the spectra of Y up by a constant.

Theorem 6. *Let X be a connected, infinite graph with finite maximum degree. Given $\epsilon > 0$, there exists $c = c(X, \epsilon) > 0$, such that for every $Y \in \mathcal{C}(X)$,*

$$|\{\lambda \in \text{spectrum of } Y : \lambda \geq \rho(X) - \epsilon\}| \geq c|V(Y)|.$$

Proof. Let $Y \in \mathcal{C}(X)$ with eigenvalues $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$ and cover map $f : V(X) \rightarrow V(Y)$. Given $\epsilon > 0$, let $m = |\{i : \lambda_i \geq \rho(X) - \epsilon\}|$.

From (2) we know that $\rho(X) = \limsup_{s \rightarrow +\infty} \sqrt[2s]{t_{2s}(x)}$, for any vertex $x \in V(X)$. Since $\rho(X) > 0$, it follows that there exists an integer $N = N(X, \epsilon) > 1$ such that $\rho(X) > \frac{\epsilon}{N}$. Obviously, if x and y are in the same orbit of $\text{Aut}(X)$, then $t_r(x) = t_r(y)$ for any non-negative integer r . Hence, the fact that X has finitely many automorphism orbits implies that there exists a non-negative integer $s_0 = s(X, \epsilon)$ such that $t_{2s}(x) \geq (\rho(X) - \frac{\epsilon}{N})^{2s}$, for each $s \geq s_0$ and any $x \in V(X)$.

It is easy to see that the number of closed walks of length r in X starting at vertex $x \in V(X)$ is less than or equal to the number of closed walks of length r in Y starting at vertex $f(x) \in V(Y)$. Hence, $\Phi_r(Y) \geq \sum_{\substack{y \in V(Y) \\ y=f(x)}} t_r(x)$ for each non-negative integer r . From the previous two relations it follows that $\Phi_{2s}(Y) \geq n(\rho(X) - \frac{\epsilon}{N})^{2s}$ for $s \geq s_0$.

Let K be a positive constant that does not depend on Y and is larger than $\lambda_1(Y)$. We can take $K = \Delta(X)$, the maximum degree of X , for example. From the previous inequality we deduce

$$\begin{aligned} \text{tr}(K \cdot I + A(Y))^{2l} &= \sum_{i=0}^{2l} \binom{2l}{i} K^{2l-i} \Phi_i(Y) \geq \sum_{j=s_0}^l \binom{2l}{2j} K^{2l-2j} \Phi_{2j}(Y) \\ &\geq n \sum_{j=s_0}^l \binom{2l}{2j} K^{2l-2j} \left(\rho(X) - \frac{\epsilon}{N}\right)^{2j} \\ &\geq n \sum_{j=0}^l \binom{2l}{2j} K^{2l-2j} \left(\rho(X) - \frac{\epsilon}{N}\right)^{2j} \\ &\quad - n \sum_{j=0}^{s_0-1} \binom{2l}{2j} K^{2l-2j} \left(\rho(X) - \frac{\epsilon}{N}\right)^{2j} \\ &\geq \frac{n}{2} \left(K + \rho(X) - \frac{\epsilon}{N}\right)^{2l} - ns_0 \binom{2l}{2s_0} K^{2l} \end{aligned}$$

for each $l \geq 2s_0$. Since $K \cdot I$ and $A(Y)$ commute, it follows that

$$\begin{aligned} \text{tr}(K \cdot I + A)^{2l} &= \sum_{i=1}^n (K + \lambda_i(Y))^{2l} \\ &\leq (n - m)(K + \rho(X) - \epsilon)^{2l} + m(2K)^{2l}. \end{aligned}$$

Hence, we obtain

$$\frac{m}{n} \geq \frac{\frac{(K + \rho(X) - \frac{\epsilon}{N})^{2l}}{2} - (K + \rho(X) - \epsilon)^{2l} - s_0 \binom{2l}{2s_0} K^{2l}}{(2K)^{2l} - (K + \rho(X) - \epsilon)^{2l}} \tag{3}$$

for each $l \geq 2s_0$.

Now

$$\lim_{l \rightarrow \infty} \sqrt[2l]{\frac{(K + \rho(X) - \frac{\epsilon}{N})^{2l}}{2}} = K + \rho(X) - \frac{\epsilon}{N}$$

and

$$\begin{aligned} \lim_{l \rightarrow \infty} \sqrt[2l]{2(K + \rho(X) - \epsilon)^{2l} + s_0 \binom{2l}{2s_0} K^{2l}} &= \max(K + \rho(X) - \epsilon, K) \\ &< K + \rho(X) - \frac{\epsilon}{N} \end{aligned}$$

imply that there exists $l_0 = l(X, \epsilon)$ such that

$$\frac{(K + \rho(X) - \frac{\epsilon}{N})^{2l}}{2} - (K + \rho(X) - \epsilon)^{2l} - s_0 \binom{2l}{2s_0} K^{2l} > (K + \rho(X) - \epsilon)^{2l}$$

for each $l \geq l_0$. Hence,

$$\frac{m}{n} > \frac{(K + \rho(X) - \epsilon)^{2l_0}}{(2K)^{2l_0} - (K + \rho(X) - \epsilon)^{2l_0}} = c(X, \epsilon) > 0. \quad \square$$

By using a similar argument as before, we can also prove a similar result to Theorem 6 for the smallest eigenvalues. Note that we need an extra hypothesis since there are classes of graphs that have eigenvalues bounded from below by a constant. For examples, line graphs have all their eigenvalues at least -2 .

Theorem 7. *Let X be a connected, infinite graph with finite maximum degree. Given $\epsilon > 0$, there exist a non-negative integer $g = g(X, \epsilon)$ and $c = c(X, \epsilon) > 0$, such that for every graph $Y \in \mathcal{C}(X)$ with no odd cycles of length less than g ,*

$$|\{\mu \in \text{spectrum of } Y : \mu \leq -(\rho(X) - \epsilon)\}| \geq c|V(Y)|.$$

Proof. Let $Y \in \mathcal{C}(X)$ with eigenvalues $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$ and cover map $f : V(X) \rightarrow V(Y)$. Given $\epsilon > 0$, let $m = |\{i : \lambda_i \leq -(\rho(X) - \epsilon)\}|$.

The proof continues now similarly to the proof of Theorem 6. There exist $N = N(X, \epsilon) > 1$, $s_0 = s(X, \epsilon)$ and $l_0 = l(X, \epsilon)$ with $l_0 \geq 2s_0$ such that

$$\frac{(K + \rho(X) - \frac{\epsilon}{N})^{2l_0}}{2} - (K + \rho(X) - \epsilon)^{2l_0} - s_0 \binom{2l_0}{2s_0} K^{2l_0} > (K + \rho(X) - \epsilon)^{2l_0}.$$

Consider

$$\text{tr}(K \cdot I - A(Y))^{2l} = \sum_{i=1}^n (K - \lambda_i)^{2l} \leq (n - m)(K + \rho(X) - \epsilon)^{2l} + m(2K)^{2l}.$$

Let $g(X, \epsilon) = 2l_0$. If Y has no odd cycles of length less than $2l_0$, then

$$\begin{aligned} \text{tr}(K \cdot I - A(Y))^{2l_0} &= \sum_{j=0}^{l_0} \binom{2l_0}{2j} K^{2l_0-2j} \Phi_{2j}(Y) \\ &\geq \sum_{j=s_0+1}^{l_0} \binom{2l_0}{2j} K^{2l_0-2j} \Phi_{2j}(Y) \\ &\geq n \sum_{j=0}^{2l_0} \binom{2l_0}{2j} K^{2l_0-2j} \left(\rho(X) - \frac{\epsilon}{N}\right)^{2j} - s_0 \binom{2l_0}{2s_0} K^{2l_0} \\ &\geq \frac{n}{2} \left(K + \rho(X) - \frac{\epsilon}{N}\right)^{2l_0} - s_0 \binom{2l_0}{2s_0} K^{2l_0}. \end{aligned}$$

From the previous two inequalities, we deduce

$$\begin{aligned} \frac{m}{n} &\geq \frac{\frac{(K + \rho(X) - \frac{\epsilon}{N})^{2l_0}}{2} - (K + \rho(X) - \epsilon)^{2l_0} - s_0 \binom{2l_0}{2s_0} K^{2l_0}}{(2K)^{2l_0} - (K + \rho(X) - \epsilon)^{2l_0}} \\ &> \frac{(K + \rho(X) - \epsilon)^{2l_0}}{(2K)^{2l_0} - (K + \rho(X) - \epsilon)^{2l_0}}. \end{aligned}$$

This proves the theorem. \square

In his Ph.D. thesis [9], Greenberg also introduced the notion of Ramanujan graph for general finite graphs (not necessarily regular). See also [14] for more details. A finite graph Y is called *Ramanujan* if for any eigenvalue $\lambda \neq \pm\chi(\tilde{Y})$ of Y , the inequality $|\lambda| \leq \rho(\tilde{Y})$ holds. If Y is regular, then we obtain the definition given by Lubotzky et al. in [16] where an infinite sequence of regular Ramanujan graphs is constructed when the degree equals $p + 1$, $p \equiv 1 \pmod 4$ prime number. In [4], this construction is extended and an infinite sequence of regular Ramanujan graphs is produced for the degree equal to a prime plus one. We are not aware of any similar results for irregular graphs as of yet.

Friedman [8] has recently proved that almost all regular graphs are almost Ramanujan. For similar results regarding irregular graphs, see [7]. Hoory [10] proved that if Y is a finite graph with average degree d , then $\rho(\tilde{Y}) \geq 2\sqrt{d - 1}$. Hoory used this result to prove a generalization of the asymptotic Alon–Boppana theorem. Denote by $B_r(v)$ the ball of radius r around v . A graph Y has an r -robust average-degree d if for every vertex v the graph induced on $V(Y) \setminus B_r(v)$ has average degree at least d . Hoory’s generalization is the following result.

Theorem 8. *Let Y_i be a sequence of graphs such that Y_i has an r_i -robust average degree $d \geq 2$, where $\lim_{i \rightarrow +\infty} r_i = +\infty$. Then*

$$\liminf_{i \rightarrow \infty} \lambda(Y_i) \geq 2\sqrt{d - 1}.$$

Acknowledgments

I am grateful to my Ph.D. advisors: David Gregory, Ram Murty and David Wehlau for their support and encouragement. I thank Shlomo Hoory for translating and adapting Greenberg's argument from Section 3 and an anonymous referee for his(her) comments.

References

- [1] N. Alon, Eigenvalues and Expanders, *Combinatorica* 6 (1986) 83–96.
- [2] S.M. Cioabă, On the extreme eigenvalues of regular graphs, *Journal of Combinatorial Theory, Series B*, in press.
- [3] S.M. Cioabă, Eigenvalues, expanders and gaps between primes, Ph.D. thesis, Queen's University at Kingston, Ont., Canada, 2005.
- [4] G. Davidoff, P. Sarnak, A. Vallette, *Elementary Number Theory, Group Theory and Ramanujan Graphs*, Cambridge University Press, 2003.
- [5] K. Feng, W.-C. Winnie Li, Spectra of hypergraphs and applications, *J. Number Theory* 60 (1) (1996) 1–22.
- [6] J. Friedman, Some geometric aspects of graphs and their eigenfunctions, *Duke Math. J.* 69 (1993) 487–525.
- [7] J. Friedman, Relative expanders or weakly relatively Ramanujan graphs, *Duke Math. J.* 118 (2003) 19–35.
- [8] J. Friedman, A proof of Alon's second eigenvalue conjecture, *Memoirs AMS*, in press.
- [9] Y. Greenberg, Spectra of graphs and their covering trees, Ph.D. thesis, Hebrew University of Jerusalem, 1995 [in Hebrew].
- [10] S. Hoory, A lower bound on the spectral radius of the universal cover of a graph, *J. Combin. Theory Ser. B* 93 (2005) 33–43.
- [11] W.-C. Winnie Li, *Number Theory with Applications*, Series of University Mathematics, World Scientific, 1996.
- [12] F.T. Leighton, Finite common coverings of graphs, *J. Combin. Theory Ser. B* 33 (1982) 231–238.
- [13] A. Lubotzky, *Discrete Groups, Expanding Graphs and Invariant Measures*, Progress in Mathematics, Birkhäuser, Basel, Boston, Berlin, 1994.
- [14] A. Lubotzky, Cayley graphs: eigenvalues, expanders and random walks, surveys in combinatorics, *London Math. Soc. Lecture Note Ser.* 218 (1995) 155–189.
- [15] A. Lubotzky, T. Nagnibeda, Not every uniform tree covers Ramanujan graphs, *J. Combin. Theory Ser. B* 74 (1998) 202–212.
- [16] A. Lubotzky, R. Phillips, P. Sarnak, Ramanujan graphs, *Combinatorica* 8 (3) (1988) 261–277.
- [17] J.-P. Serre, Répartition asymptotique des valeurs propres de l'opérateur de Hecke T_p , *J. Amer. Math. Soc.* 10 (1) (1997) 75–102.
- [18] D.B. West, *Introduction to Graph Theory*, second ed., Prentice Hall, New Jersey, 2001.